



Multisociety Gluteal Fat Grafting Task Force Issues Safety Advisory Urging Practitioners to Reevaluate Technique

Date Issued: January 31, 2018

Dear Colleagues,

An Inter-Society Gluteal Fat Grafting Task Force** has analyzed deaths from gluteal fat injection ("Brazilian Butt Lift" or "BBL") and offers the following advisory statement: The death rate of approximately 1/3000 is the highest for any aesthetic procedure. In 2017, there were three deaths in the state of Florida alone. Every surgeon performing BBLs should immediately reevaluate his or her technique.

Some patients have died when their surgeon said they had injected into the subcutaneous fat layer, but all autopsies of deceased BBL patients have had these findings in common: 1) fat in the gluteal muscles; 2) fat beneath the muscles; 3) damage to the superior or inferior gluteal vein; 4) massive fat emboli in the heart and/or lungs. No postmortem has yet shown a case of death with fat only in the subcutaneous space; this means that surgeons have injected more deeply than they had intended. The mechanism of death is presumed to be high pressure extravascular grafted fat entering the circulation via tears in the large gluteal veins with subsequent embolization to the heart and lungs.

The task force, therefore, offers these suggestions*:

1. Stay as far away from the gluteal veins and sciatic nerve as possible. Fat should only be grafted into the superficial planes, with the subcutaneous space considered safest. If the aesthetic goal requires more fat than can be placed in the subcutaneous layer the surgeon should consider staging the procedure rather than injecting deep.
2. Concentrate on the position of the cannula tip throughout every stroke to assure there is no unintended deeper pass, particularly in the medial half of the buttock overlying the critical structures.

3. Use access incisions that best allow a superficial trajectory for each part of the buttock; avoid deep angulation of the cannula; and palpate externally with the non-dominant hand to assure the cannula tip remains superficial.
4. Use instrumentation that offers control of the cannula; avoid bendable cannulas and mobile luer connections. Vibrating injection cannulas may provide additional tactile feedback.
5. Injection should only be done while the cannula is in motion in order to avoid high pressure bolus injections.
6. The risk of death should be discussed with every prospective BBL patient.
7. These are links to three helpful articles:
 - [Staying Safe During Gluteal Fat Transplantation, Plastic and Reconstructive Surgery, January 2018](#)
 - [Report on Mortality from Gluteal Fat Grafting: Recommendations from the ASERF Task Force; Aesthetic Surgery Journal, July 2017](#)
 - [Acceptable Risk: Who Decides? Editorial by Foad Nahai, MD, Aesthetic Surgery Journal, July 2017](#)

Research projects overseen by the task force and funded by The Plastic Surgery Foundation (PSF) and the Aesthetic Surgery Education and Research Foundation (ASERF) are underway. They will correlate deep and topographical anatomy, define danger zones, and try to understand the mechanism of embolization. The ability to safely perform this procedure in the future is dependent upon this research. Every surgeon performing BBL should click on one of these links to make their contribution.


Members of the task force have also assisted coroners during autopsies, and this has provided invaluable safety information. If you become aware of a fatality, immediately contact the task force co-chairs care of Keith Hume, executive director of The PSF, at khume@plasticsurgery.org.

Your societies will keep you updated with all developments.


Sincerely,



Dan Mills, MD
Gluteal Fat Grafting
Task Force co-chair



J. Peter Rubin, MD
Gluteal Fat Grafting
Task Force co-chair



Renato Saltz, MD
Gluteal Fat Grafting
Task Force co-chair

* The information in this Advisory Statement while setting forth the strong recommendations of the Task Force, should not be considered inclusive of all methods of properly performing buttock augmentation with fat transfer or as a statement of the standard of care or as a mandate to strictly follow the recommendations of the Task Force. This Advisory Statement is not intended to substitute for the independent professional judgment of the treating plastic surgeon nor for the individual variation among patients. The Members of the Multi-Society Task Force and the participating societies assume no responsibility or liability for injury arising out of any use of the information contained in this Advisory Statement.

** The Inter-Society Gluteal Fat Grafting Task Force represents leading clinical plastic surgery societies, including the American Society of Plastic Surgeons (ASPS), the American Society for Aesthetic Plastic Surgery (ASAPS), and the International Society of Aesthetic Plastic Surgeons (ISAPS). Additionally, two scientific societies, the International Society of Plastic & Regenerative Surgeons (ISPRES) and the International Federation for Adipose Therapeutics and Science (IFATS) are represented and provide scientific support. The efforts of the Task Force build upon a foundation of important work by the Aesthetic Surgery Education and Research Foundation (ASERF), the American Society of Plastic Surgeons (ASPS) Regenerative Medicine Committee, and the International Society of Aesthetic Plastic Surgery (ISAPS) Patient Safety Committee. The Task Force is an unprecedented collaborative effort to address a major patient safety concern, investigate factors that lead to increased risk with gluteal fat grafting, perform scientific studies to improve safety, and educate plastic surgeons.